

# Columbia-Suicide Severity Rating Scale (C-SSRS)

Screen with Triage Points for Primary Care

Ask questions that are in bold and underlined.

Past month

Ask Questions 1 and 2	YES	NO
1) <b><u>Have you wished you were dead or wished you could go to sleep and not wake up?</u></b>		
2) <b><u>Have you had any actual thoughts of killing yourself?</u></b>		
If YES to 2, ask questions 3, 4, 5, and 6. If NO to 2, go directly to question 6.		
3) <b><u>Have you been thinking about how you might do this?</u></b> e.g. "I thought about taking an overdose but I never made a specific plan as to when where or how I would actually do it...and I would never go through with it."		
4) <b><u>Have you had these thoughts and had some intention of acting on them?</u></b> as opposed to "I have the thoughts but I definitely will not do anything about them."		
5) <b><u>Have you started to work out or worked out the details of how to kill yourself?</u></b> Do you intend to carry out this plan?		
6) <b><u>Have you ever done anything, started to do anything, or prepared to do anything to end your life?</u></b> Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.  If YES, ask: <b><u>Was this within the past 3 months?</u></b>	<b>Lifetime</b>	
	<b>Past 3 Months</b>	

## Response Protocol to C-SSRS Screening (use protocol in accordance with clinical judgment)

Risk Level	Suggested Interventions
<p style="text-align: center;"><b>High Risk</b></p> <p>Suicidal ideation with intent or intent with plan in past month (C-SSRS Suicidal Ideation #4 or #5) <i>or</i> Suicidal behavior within past three months (C-SSRS Suicidal Behavior)</p>	<p>Call 911 for transport to the emergency room or contact community crisis line in your area to provide on-site evaluation.</p> <p>Place individual in a room that is away from exits but close to staff where patient is observed at all times until help arrives.</p>
<p style="text-align: center;"><b>Medium Risk</b></p> <p>Suicidal ideation WITHOUT plan, intent or behavior in past month (C-SSRS screen #2 or #3) <i>or</i> Suicidal behavior more than three months ago (C-SSRS Suicidal Behavior)</p>	<p>If patient is already receiving mental health treatment, get release of information. If not, refer to mental health provider for further assessment (within one week).</p> <p>Consider pharmacological treatment.</p> <p>Provide education on safe fire arms storage, suicide warning signs and 1-800-273-TALK (8255) and local contacts.</p>
<p style="text-align: center;"><b>Low Risk</b></p> <p>Wish to die (C-SSRS Suicidal Ideation #1) without plan, intent or behavior <i>or</i> Suicidal ideation more than one month ago WITHOUT plan, intent or behavior (C-SSRS screen #2 or #3)</p>	<p>Assess for any other mental health or substance use conditions and consider behavioral health and/or pharmacological treatment.</p> <p>Provide education on safe fire arms storage, suicide warning signs and 1-800-273-TALK (8255) and local contacts.</p>

Ensure that you have a clear and simple office protocol in place for patients who are suicidal. Explore the following resource for guidance <http://www.sprc.org/sites/default/files/OfficeProtocolDevelopmentGuide.pdf>